

ACCESSORY NAVICULAR SYNDROME



American College of
Foot and Ankle Surgeons

What is the Accessory Navicular?

The accessory navicular (os navicularum or os tibiale externum) is an extra bone or piece of cartilage located on the inner side of the foot just above the arch. It is incorporated within the posterior tibial tendon, which attaches in this area.



An accessory navicular is congenital (present at birth). It is not part of normal bone structure and therefore is not present in most people.

What is Accessory Navicular Syndrome?

People who have an accessory navicular often are unaware of the condition if it causes no problems. However, some people with this extra bone develop a painful condition known as accessory navicular syndrome when the bone and/or posterior tibial tendon are aggravated. This can result from any of the following:

- Trauma, as in a foot or ankle sprain
- Chronic irritation from shoes or other footwear rubbing against the extra bone
- Excessive activity or overuse

Many people with accessory navicular syndrome also have flat feet (fallen arches). Having a flat foot puts more strain on the posterior tibial tendon, which can produce inflammation or irritation of the accessory navicular.



Signs and Symptoms of Accessory Navicular Syndrome

Adolescence is a common time for the symptoms to first appear. This is a time when bones are maturing and cartilage is developing into bone. Sometimes, however, the symptoms do not occur until adulthood. The signs and symptoms of accessory navicular syndrome include:

- A visible bony prominence on the midfoot (the inner side of the foot, just above the arch)
- Redness and swelling of the bony prominence
- Vague pain or throbbing in the midfoot and arch, usually occurring during or after periods of activity

Diagnosis

To diagnose accessory navicular syndrome, the foot and ankle surgeon will ask about symptoms and examine the foot, looking for skin irritation or swelling. The doctor may press on the bony prominence to assess the area for discomfort. Foot structure, muscle strength, joint motion, and the way the patient walks may also be evaluated.

X-rays are usually ordered to confirm the diagnosis. If there is ongoing pain or inflammation, an MRI or other advanced imaging tests may be used to further evaluate the condition.

Treatment: Non-Surgical Approaches

The goal of non-surgical treatment for accessory navicular syndrome is to relieve the symptoms. The following may be used:

- **Immobilization.** Placing the foot in a cast or removable walking boot allows the affected area to rest and decreases the inflammation.
- **Ice.** To reduce swelling, a bag of ice covered with a thin towel is applied to the affected area. Do not put ice directly on the skin.

- **Medications.** Oral nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may be prescribed. In some cases, oral or injected steroid medications may be used in combination with immobilization to reduce pain and inflammation.
- **Physical therapy.** Physical therapy may be prescribed, including exercises and treatments to strengthen the muscles and decrease inflammation. The

exercises may also help prevent recurrence of the symptoms.

- **Orthotic devices.** Custom orthotic devices that fit into the shoe provide support for the arch, and may play a role in preventing future symptoms.

Even after successful treatment, the symptoms of accessory navicular syndrome sometimes reappear. When this happens, non-surgical approaches are usually repeated.

When is Surgery Needed?

If non-surgical treatment fails to relieve the symptoms of accessory navicular syndrome, surgery may be appropriate. Surgery may involve removing the accessory bone, reshaping the area, and repairing the posterior tibial tendon to improve its function. This extra bone is not needed for normal foot function. ▲



16841 N. 31st Ave | Suite 134 | Phoenix, Arizona 85053
 Office: (623) 322-5501
 Fax: (623) 322-8996



**American College of
Foot and Ankle Surgeons**

This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of 6,200 foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.

Copyright © 2008, American College of Foot and Ankle Surgeons • www.FootPhysicians.com